


GEBCO (1979) AND IBCAO (2000) MAPS

SELECTIVE COMPARISONS



Martin Jakobsson, Stockholm University

Ron Macnab, Geological Survey of Canada

Members of the Editorial Board, IBCAO

SELECTIVE COMPARISONS OF GEBCO (1979) AND IBCAO (2000) MAPS

Martin Jakobsson, Stockholm University
martin.jakobsson@geo.su.se

Ron Macnab, Geological Survey of Canada
macnab@agc.bio.ns.ca

Members of the Editorial Board, IBCAO
arctic-bathy@ldeo.columbia.edu

Introduction

For over two decades Sheet 5.17 of the General Bathymetric Chart of the Oceans (GEBCO, Canadian Hydrographic Service, 1979) served as an authoritative portrayal of the seafloor north of 64°N. While this contour map provided a general description of major features of the seabed, evidence was accreting to indicate that many of the smaller and scientifically significant features were poorly or wrongly defined.

The International Bathymetric Chart of the Arctic Ocean (IBCAO) was developed from an accumulation of bathymetric measurements collected during past and modern expeditions. The undertaking has produced a 2.5 x 2.5 km grid and derivative maps (Jakobsson *et al.*, 2000) that represent a considerable improvement over previous descriptions of the morphology of the Arctic seabed. The most substantial improvement is in the central Arctic Ocean, where a nearly empty database was significantly enriched by the addition of historic and modern observations collected by US and British submarines (DeLaca and Gossett, 1996; Showstack, 1997), by Swedish and German icebreakers, and by depths derived from a new contour map prepared by the Russian Navy (Head Department of Navigation and Hydrography *et al.*, 1999). The database for the shallow shelves of the Laptev and East Siberian Seas has been enhanced as well, absorbing a large quantity of depth values that were extracted from a suite of small scale navigational charts issued by the Russian Navy.

This note provides examples of the above improvements by contrasting the GEBCO and IBCAO representations of the seafloor in four regions of the Arctic Ocean: the Gakkel Ridge, the Lomonosov Ridge, the Chukchi Borderland, and the continental shelves of the Chukchi, Eastern Siberian and Laptev Seas. These regions are shown on the cover page.

The GEBCO information that was used in this comparison was actually taken from a digital rendition of GEBCO Sheet 5.17, constructed from bathymetric contours that were distributed on CD-ROM as part of the GEBCO Digital Atlas (Jones *et al.*, 1994). The contours were used to develop a Cartesian depth grid (5 x 5 km grid cells) on a polar stereographic map projection with the true scale at 75°N (Geological Survey of Canada, 1994). These map parameters are consistent with the original paper map, and they are also compatible with those of the new IBCAO map. Both the GEBCO and the IBCAO grids were created by means of the GMT program *surface* (Smith and Wessel, 1990). The comparisons that follow consist of side-by-side shaded relief portrayals of three study areas, constructed from the GEBCO and IBCAO grids using identical illumination parameters and color tables (Figure 1).



Figure 1. The color table used to create the shaded relief maps in figures 2-5.

The Gakkel Ridge

The Gakkel Ridge is a spreading ridge that separates the Nansen and Amundsen Basins. It is a continuation of the mid-oceanic ridge system into the Arctic Ocean (e.g., Johnson and Heezen, 1967; Vogt, 1986).

The GEBCO map portrays the western end of the Gakkel Ridge with a smooth and highly generalized morphology (Figure 2). On the IBCAO map, the Ridge flanks feature a blocky configuration, while the Ridge's axial valley is continuous, more pronounced, and deeper, reaching depths below 5000 m and containing several inter-axial highs.




Figure 2. A comparison of the Gakkel Ridge, portrayed by IBCAO (left) and GEBCO (right).

The Lomonosov Ridge

Measuring about 1700 km in length, the Lomonosov Ridge is considered to be of continental origin, a sliver that was separated from the Kara and Barents shelves and transported to its present position by sea-floor spreading (e.g., Sweeney *et al.*, 1982; Kristoffersen, 1990).

On the GEBCO map, the Lomonosov Ridge is a continuous feature that extends from the continental shelf off Ellesmere Island towards the North Pole, where it changes direction

slightly and continues along the 140°E meridian to the continental shelf off the New Siberian Islands. The most striking discrepancies between the GEBCO and IBCAO portrayals occur between the North Pole and the Siberian continental shelf (Figure 3). The new model shows a far more complex morphology, with a ridge that is broken into several smaller segments, and a pronounced kink that projects into Amundsen Basin. Moreover, the crest is generally shallower, wider, and notably flat-topped at several sections.




Figure 3. A comparison of the Lomonosov Ridge in the vicinity of the North Pole, portrayed by IBCAO (left) and GEBCO (right).

The Chukchi Borderland

Measuring about 600 by 700 km, the Chukchi Borderland extends from the continental shelves of Eastern Siberia and Western Alaska into the deep Amerasian Basin. It contains three north-south trending structures: Northwind Ridge, Chukchi Cap and Arlis Plateau - the last feature is commonly considered to be a part of the Mendeleev Ridge. Adjacent to these ridge-like structures lie the Northwind, Chukchi and Mendeleev Abyssal Plains.

GEBCO and IBCAO comparisons are limited here to the Northwind Ridge and the Chukchi Cap. Both are among the more extensively mapped features in the Arctic Ocean, see e.g. the compilation map published by the US Naval Research Laboratory (Perry et al., 1985).




Figure 4. A comparison of the Northwind Ridge and the Chukchi Cap in the Chukchi Borderland, portrayed by IBCAO (left) and GEBCO (right)

On the IBCAO map, these features appear to be more highly segmented than on previous maps, and their surfaces display irregularities that suggest complex sea-floor processes (Figure 4).

The continental shelves of the Eastern Siberian and Laptev Seas

The shelf areas of the Eastern Siberian and Laptev Seas are shallow, generally less than 50 m deep, with a very flat relief.

With a 50 m minimum depth contour, the GEBCO map reveals very little bathymetric information over these huge shelf areas. On the other hand, a shaded relief implementation of the IBCAO map is effective at portraying depth variations of only tens of metres, accentuating the shallow paleo-river valleys that incise these areas, along with low-relief erosional and depositional features (Figure 5).




Figure 5. Oblique view of the continental shelves of the Eastern Siberian and Laptev Seas. The coloration of shallow regions is not the same as in Figures 1-4 on account of a different shading technique that was applied to emphasize low-relief features.

Members of the Editorial Board for IBCAO

Harald BREKKE	Harald.Brekke@npd.no	Norway
Norwegian Petroleum Directorate		
PO Box 600		
4001 Stavanger		
Norway		

Norman Z. CHERKIS	cherkis@excite.com	USA
Five Oceans Consultants		
6300 Saddle Tree Drive		
Alexandria VA DC 22310-2915		
USA		

Bernie COAKLEY	bcoakle@mailhost.tcs.tulane.edu	USA
Department of Geology		
Tulane University		
New Orleans LA 70118		
USA		

David DIVINS National Geophysical Data Center E/GC3 325 Broadwat Boulder CO 80303 USA	ddivins@ngdc.noaa.gov	USA
Valeriy FOMCHENKO Head Department of Navigation and Oceanography Russian Federation Navy 8,11 Liniya, B-34 199034 St. Petersburg Russia	gunio@g-ocean.spb.su	Russia
Garrik GRIKUROV VNIOkeangeologia 1 Anglisky Avenue 190121 St. Petersburg Russia	grikurov@mail.lanck.net	Russia, IASC
Neil R. GUY International Hydrographic Bureau BP 445 MC 98011 Monaco Cedex Principauté de Monaco	dir1@ihb.mc	IHO
Hilmar HELGASON Icelandic Hydrographic Service Seljaveg 32 127 Reykjavik Iceland	hilmar@lhg.is	Iceland
Martin JAKOBSSON Stockholm University Department of Geology and Geochemistry S-10691 Stockholm Sweden	martin.jakobsson@geo.su.se	Sweden
Ron MACNAB (Chairman) Geological Survey of Canada PO Box 1006 Dartmouth NS B2Y 4A2 Canada	macnab@agc.bio.ns.ca	Canada
Sergei MASCHENKOV VNIOkeangeologia 1 Anglisky Avenue 190121 St. Petersburg Russia	mascha@vniio.nw.ru	Russia

Hans-Werner SCHENKE schenke@awi-bremerhaven.de Germany
Alfred Wegener Institute
Postfach 120161
D-27515 Bremerhaven
Germany

Dmitri TRAVIN d.travin@unesco.org IOC
Intergovernmental Oceanographic Commission
1, rue Miollis
75732 Paris, Cedex 15
France

John WOODWARD jjw@fomfrv.dk Denmark
Royal Danish Administration of Navigation and Hydrography
Overgaden o. Vandet 62B
DK-1023 Copenhagen K
Denmark

Bibliography

Canadian Hydrographic Service. *General Bathymetric Chart of the Oceans (GEBCO) Sheet 5.17*; map 1:6,000,000. Canadian Hydrographic Service, Ottawa, 1979.

DeLaca, T., and J. Gossett. *Nuclear Submarines and Oceanography in the Arctic*; Witness the Arctic: Chronicles of the Arctic System Science Research Program, Arctic Research Consortium of the United States, Spring 1996, Vol. 3, No. 1., 1996.

Geological Survey of Canada. *Bathymetric and topographic shaded relief north of 64N*; polar stereographic projection, variable scale, distributed in hardcopy or digital form. GSC Open File 2900, Geological Survey of Canada, Dartmouth NS, 1994.

Head Department of Navigation and Oceanography, All-Russia Research Institute for Geology and Mineral Resources of the World Ocean, and Russian Academy of Sciences. *Bottom relief of the Arctic Ocean*; map 1:5 Million. Head Department of Navigation and Oceanography, St. Petersburg, 1999.

Jakobsson, M., N.Z. Cherkis, J. Woodward, R. Macnab, and B. Coakley. *New grid of Arctic bathymetry aids scientists and mapmakers*; Eos, Transactions, American Geophysical Union, in press.

Jones, M.T., A.R. Tabor, and P. Weatherall. *GEBCO Digital Atlas: CD-ROM and supporting Volume*. British Oceanographic Data Centre, Birkenhead, UK., 1994.

Johnson, G. L., and Heezen, B. C. *The Arctic Mid-Oceanic Ridge*; Nature, vol. 215, pp. 724-725, 1967.

Kristoffersen, Y. *Eurasia Basin*; In: A. Grantz, L. Johnson, and F.J. Sweeney (Eds.), *The Arctic Ocean Region*; DNAG, vol. L: pp. 365-378, 1990.

Perry, R.K., H.S. Fleming, J.R. Weber, Y. Kristofferson, J.K. Hall, A. Grantz, G.L. Johnson. *Bathymetry of the Arctic Ocean*; map 1:4,704,075. Naval Research Laboratory, Washington, 1985.

Showstack, R., *Navy to Release Storehouse of Arctic Ocean Data*. EOS, Trans. Amer. Geophys. Un. vol. 78, no. 35, pp. 370-371, 1997.

Smith, W. H. F., and P. Wessel, *Gridding with continuous curvature splines in tension*, Geophysics, vol. 55, pp. 293-305, 1990.

Sweeney, J. F., R. R. Weber, , and S. M. Blasco. *Continental ridges in the Arctic Ocean: LOREX constraints*; Tectonophysics, v. 89, p. 217-238, 1982.

United States Arctic Research Commission. *US Arctic Research Commission and US Navy Announce Release of Arctic Ocean Bathymetry Data*; Press Release, August 21, 1997, US Arctic Research Commission, Arlington VA, USA.

Vogt, P. R. *Geophysical and geochemical signatures and plate tectonics*; In: The Nordic Seas, Hurdle, B. G. (ed.), Springer-Verlag, pp. 413-662, 1986.